

Synergism between *Phyllobacterium* sp. (N₂-fixer) and *Bacillus licheniformis* (P-solubilizer), both from a semiarid mangrove rhizosphere

Adriana Rojas^a, Gina Holguin^{a,b}, Bernard R. Glick^b, Yoav Bashan^{a,*}

^a Environmental Microbiology, The Center for Biological Research of the Northwest (CIB), A.P. 128, La Paz, Baja California Sur 23000, Mexico

^b Department of Biology, University of Waterloo, Waterloo, Ont., Canada N2L 3G1

Received 30 July 2000; received in revised form 22 December 2000; accepted 26 December 2000

Abstract

Mangrove seedlings were treated with a mixture of two bacterial species, the slow-growing, N₂-fixing bacterium *Phyllobacterium* sp. and the fast-growing, phosphate-solubilizing bacterium *Bacillus licheniformis*, both isolated from the rhizosphere from black, white, and red mangroves of a semiarid zone. Nitrogen fixation and phosphate solubilization increased when the mixture was used compared to the effects observed when adding individual cultures, notwithstanding that there was no increase in bacterial multiplication under these conditions. Inoculation of black mangrove seedlings in artificial seawater showed the mixture performed somewhat better than inoculation of the individual bacterium; more leaves were developed and higher levels of ¹⁵N were incorporated into the leaves, although the total nitrogen level decreased. This study demonstrates that interactions between individual components of the rhizosphere of mangroves should be considered when evaluating these bacteria as plant growth promoters. © 2001 Federation of European Microbiological Societies. Published by Elsevier Science B.V. All rights reserved.

Keywords: Mangrove; Mutualism; Nitrogen fixation; Phosphate solubilization; Plant growth-promoting bacterium; *Phyllobacterium*; *Bacillus licheniformis*

1. Introduction

Sediment and rhizosphere microorganisms are the major biological components that contribute to the productivity of mangroves [1–6]. Because of diverse microbial activity, mangrove ecosystems are one of the three most productive ecosystems known, along with rain forests and coral reefs [7]. Yet, mangroves are alarmingly and systematically being deforested, similarly to rain forests [8,9]. To aid in the reforestation of mangroves, inoculation of the seedlings with plant growth-promoting bacteria (PGPBs) has been suggested [10,11,12], similarly to what has been done in agriculture [13–16] and temperate forestry [17].

At the present time, little is known about the PGPBs and their mechanisms and interactions that are normally associated with mangroves. Previous work on PGPBs that interact with mangroves includes the observation that the diazotrophic cyanobacteria *Microcoleus chthonoplastes* improved N₂ fixation [12] and nitrogen incorporation in

black mangrove seedlings [10]. In addition, the terrestrial halotolerant bacterium *Azospirillum halopraeferens* [18] and halotolerant *Azospirillum brasilense* Cd [19] can successfully colonize black mangrove roots in seawater [11]. More recently, several potential PGPBs of mangrove origin (i.e. *Bacillus licheniformis*, *M. chthonoplastes*, *Phyllobacterium myrsinacearum*, *Vibrio aestuarianus*, and *Vibrio proteolyticus* [12,20]) were shown to promote the growth of the annual, potential oilseed seaweed *Salicornia bigelovii*, which shares the semiarid mangrove ecosystem with the trees [21].

In agricultural and forestry inoculation practices, mixing of two or more microbial species often has a more positive effect on plant growth than the use of a single bacterium [22–28].

The interaction of N₂-fixing bacteria with other bacteria can inhibit or promote their diazotrophic activity [29,30]. The degradation of cellulose by *Cellulomonas* sp. provided *Azospirillum* sp. with a usable carbon source to obtain energy for N₂ fixation. The contribution of *Azospirillum* to *Cellulomonas* is fixed nitrogen [31]. The association between different *Azospirillum* species and N₂-fixer *Bacillus polymyxa* enhanced the N₂-fixing activity of the mixed

* Corresponding author. Fax: +52 (112) 54710 or 53625;
E-mail: bashan@cibnor.mx

culture compared to the pure culture of each bacterium. *Azospirillum* benefited from the products that are released from the degradation of pectin by *B. polymyxa* [32]. The N₂-fixing activity of *A. brasilense* Cd increased significantly when grown in mixed culture with the mangrove rhizosphere bacterium *Staphylococcus* sp., releasing aspartic and succinic acids into the growth medium [19]. Apart from the increase in N₂ fixation when the mangrove N₂-fixer *Listonella anguillarum* was grown together with *Staphylococcus* sp. [33], other interactions of potential mangrove PGPBs among themselves and their effect on plant growth are unknown.

The aim of this study was to explore the possible interactions in vitro of two potential mangrove PGPBs; the N₂-fixing *Phyllobacterium* sp. and the phosphate-solubilizing bacterium *B. licheniformis*.

2. Materials and methods

2.1. Organisms and growth conditions

Phyllobacterium sp. was isolated according to Holguin et al. [33] from the rhizosphere of semiarid black (*Avicennia germinans*), white (*Laguncularia racemosa*), and red (*Rhizophora mangle*) mangrove seedlings from Laguna de Balandra, Baja California Sur, Mexico. Its nitrogen-fixing capacity was assessed as described by Holguin et al. [33]. This bacterium was characterized using both fatty acid methyl ester analysis and 16S rRNA by a commercial service (Acculab, Newark, DE, USA). Although the analyses indicate this bacteria is probably *Phyllobacterium rubiacearum* or *P. myrsinacearum*, it did not produce nodules on leaves, fix atmospheric nitrogen, and its habitat is seawater. These characteristics do not fit the species description, but it did show star clusters, and its colonies and cellular characteristics do fit the genus description [34,35]. Therefore, we chose to cite this strain as *Phyllobacterium* sp. *B. licheniformis*, originally isolated from black and white mangroves with a phosphate solubilization capacity of 325 mg P l⁻¹, was obtained from the CIB culture collection. It was purified and characterized in a previous study [20]. Both of the above bacterial species were maintained for routine use in 30% glycerol (v/v) and stored in modified SRSM2 medium for *B. licheniformis* [20] and N-free medium for *Phyllobacterium* sp. at either -40 ± 3°C or in liquid nitrogen.

Black mangrove propagules were collected from Laguna de Balandra, Baja California Sur, Mexico [36,37]. They were treated and inoculated with the bacteria as previously described [10,12].

All experiments were done in vitro. Bacteria were grown in un baffled Erlenmeyer flasks on an environmentally controlled rotary shaker at 150 rpm (*B. licheniformis*), or without agitation (*Phyllobacterium* sp.) at 30 ± 1°C. Three culture media were used; (i) both liquid and solid

minimal medium SRSM2 [20] supplemented with 1.1 g K₂HPO₄·3H₂O l⁻¹ (Sigma), (ii) N-free minimal medium for marine bacteria containing (g l⁻¹): 0.075, K₂HPO₄·3H₂O; 0.028, FeSO₄·7H₂O; 1.25, malic acid; 1.25, citric acid; 2, D-glucose; 2, D-mannitol; 1, myo-inositol; 23.4, NaCl; 1.5, KCl; 24, MgSO₄·7H₂O; 2.9, CaCl₂·2H₂O; 10 ml glycerol; (µg l⁻¹) 0.01, biotin; 0.02, pyridoxine; 0.4, Na₂MoO₄·2H₂O; 4.7, MnSO₄·H₂O; 0.56, H₃BO₃; 0.16, CuSO₄·5H₂O; 0.048, ZnSO₄·7H₂O (Holguin, G., 1997; Technical report for Consejo Nacional de Ciencia y Tecnologia, Mexico), and (iii) filtered seawater [11]. Incubation times varied according to the experiment; up to 5 days for phosphate and nitrogen fixation tests, 5 days for cultures containing mangrove seedlings for acetylene-reduction assay, and 26 days for cultures containing mangrove seedlings (7 cm tall) for total nitrogen content and ¹⁵N abundance.

Bacterial counts were done by the Plate Count Method [38] on Nutrient Broth medium (Difco) supplemented with 2% NaCl.

2.2. Analyses

Nitrogen fixation was measured by the acetylene-reduction assay using gas chromatography after 48 h of mixed incubation as shorter periods showed negligible values [33]. Root surface area was measured using the calcium nitrate gravimetric method [39]. Total nitrogen was measured by an automatic micro-Kjeldahl procedure after digestion (Digestion System 12.1009, and Kjeltec Auto 103 analyzer, Tecator, Höganäs, Sweden). Abundance of ¹⁵N in the sample was measured by Isochrom Continuous Flow, Stable Isotope Mass Spectrometer (Micromass, Manchester, UK) according to standard methods [40] and expressed as ¹⁵N in parts per thousand (δ) as previously described [10]. Phosphate solubilization was done using a modification of the method of Vazquez et al. [20] as follows: *B. licheniformis* was grown on SRSM2 solid medium for 24 h and *Phyllobacterium* sp. for 120 h on N-free medium. All colonies were aseptically scraped from the agar surface, each suspended in 1 ml of 2% NaCl, and the number of colony-forming units (cfu) ml⁻¹ determined. The two cultures were then mixed and inoculated into 25 ml of natural seawater [11] supplemented with 5 g l⁻¹ tribasic calcium phosphate (Sigma) and incubated under agitation of 150 rpm at 28°C for an additional 120 h. Three-ml samples from this culture were centrifuged at 14 000 × g for 3 min. The supernatant (2.5-ml samples) was collected and diluted to 25 ml with distilled water. Phosphate analysis was done according to standard methods of water analysis [41] as modified by the Hach Co. (CO, USA) using a Hach DR2000 spectrophotometer (Hach program # 480) at 430 nm. Light microscopy of freshly mounted cultures and spore staining (methylene blue) were done by using a Zeiss (Germany) light microscope (×1000).




Fig. 1. Growth curves of *Phyllobacterium* sp. and *B. licheniformis* in pure and mixed cultures. (A) Minimal medium SRSM2, (B) N-free minimal medium, (C) seawater. Points denoted by a different lower case letter for each bacterial growth curve separately differ significantly at $P \leq 0.05$ by one-way ANOVA. Points in each sampling time denoted by different capital letter in each subfigure separately differ significantly at $P \leq 0.05$ by the Student's *t*-test. Bars represent S.E.M. The absence of S.E.M. means that it is smaller than the point. (D) Appearance and disappearance of spores of *B. licheniformis* during growth in different culture media.




Fig. 2. (A) Nitrogen fixation (acetylene reduction) of bacterial cultures inoculated with the N₂-fixing bacteria *Phyllobacterium* sp. alone or mixed with the phosphate-solubilizing bacteria *B. licheniformis*. (B) Phosphate solubilization of *B. licheniformis* cultures alone or together with *Phyllobacterium* sp. Columns denoted by a different letter, in each subfigure, differ significantly at $P \leq 0.05$ by one-way ANOVA. Number above each column represents the bacterial population in cfu ml⁻¹. Bars represent S.E.M. Absence of S.E.M. indicates minimal S.E.M.

2.3. Experimental design and statistical analysis

All experiments were done in 10 replicates using an Erlenmeyer flask (for bacteria) or filtration flask containing one mangrove seedling as a replicate. All experiments were repeated 2–3 times. Analysis of ^{15}N was done once. Results were analyzed using one-way Analysis of Variance (ANOVA) or by the Student's *t*-test, both at $P \leq 0.05$, using Statistica software (StatSoft, Tulsa, OK, USA).

3. Results

3.1. Growth of *Phyllobacterium* sp. and *B. licheniformis* in mixed cultures

Light microscopy revealed that when the two bacterial species were grown on solid medium they formed one morphotype colony containing both species, whereas on liquid medium they grew as separate cells. Thus, the

counts on solid medium of mixed cultures done in this study could not distinguish between the different populations.

On minimal medium having a soluble P source, the fast grower *B. licheniformis* maintained its population in the first 72 h, then gradually the population decreased. The mixed population increased bacterial population in the first 24 h, and then decreased (Fig. 1A). *Phyllobacterium* sp., a slow grower, continued its growth for 120 h when grown alone in N-free medium (having a soluble P source). In mixed culture on this medium, a similar decrease in population occurred as with the previous medium (Fig. 1B). In seawater, after an initial decrease in their populations, both bacterial species increased their population over time as did the mixed population, approximately to their initial level. The decrease in the mixed population observed in the previous two media was not apparent (Fig. 1C).

Phyllobacterium sp. cells did not change their morphology in culture whether alone or in a mixture. However,




Fig. 3. Total nitrogen content of black mangrove seedlings inoculated with the N_2 -fixing bacteria *Phyllobacterium* sp. alone or mixed with the phosphate-solubilizing bacteria *B. licheniformis* (A–C). ^{15}N accumulation in the same plants (D–F). Columns denoted by a different letter, in each subfigure, differ significantly at $P \leq 0.05$ by one-way ANOVA. Bars represent S.E.M. Absence of a bar above a column indicates minimal S.E.M.

cells of the fast-growing *B. licheniformis* produced spores that appeared at different incubation times depending on the medium. In pure culture in minimal medium, endospores were observed after 72 h of incubation. In seawater, endospores were found within the first 24 h, and older cultures consisted almost exclusively of spores. A mixed culture in seawater resembled pure cultures in seawater. First there was spore formation but different from the pure culture. After 120 h there was massive reappearance of vegetative cells (Fig. 1D).

3.2. Nitrogen fixation, phosphate solubilization, and nitrogen accumulation in seedlings inoculated with mixed cultures

Acetylene reduction of mixed cultures revealed that *Phyllobacterium* sp. fixes twice the amount of nitrogen than when it is in pure culture (Fig. 2A). Similarly, phosphate solubilization by *B. licheniformis* increased significantly in mixed as compared to monocultures (Fig. 2B). Black mangrove seedlings inoculated with either pure or mixed cultures showed similar levels of nitrogen fixation (10.7 nmol ethylene per plant per 7 days). All inoculation treatments reduced the nitrogen content in the leaves, stems, and roots of the plants (Fig. 3A–C). The ^{15}N content in the plant leaves and stems was significantly higher in mixed culture than in uninoculated plants (Fig. 3D–F).

3.3. Root colonization and the effect of inoculation on black mangrove seedlings

All inoculations significantly increased the number of true leaves of black mangrove seedlings over the uninoculated control; however, there was no significant difference




Fig. 4. Effect of inoculation with the N_2 -fixing bacteria *Phyllobacterium* sp. and the phosphate-solubilizing bacteria *B. licheniformis* alone or together on the number of true leaves developed by the seedlings. Columns denoted by a different letter differ significantly at $P \leq 0.05$ by one-way ANOVA. Bars represent S.E.M.

among the inoculation treatments (Fig. 4). None of the inoculations had any effect on the root surface area of the seedlings. Root colonization of the seedlings was at a level of 2×10^4 cfu g^{-1} dw. root for both pure cultures and 1×10^5 cfu g^{-1} dw. root for the mixed culture. We could not differentiate between the two bacterial species as they grew as one mixed colony on the root surface.

4. Discussion

Semiarid mangrove forests thrive in lagoons that are lacking sources of both dissolved phosphorus and nitrogen [5,20], essential growth elements for any plant species. Nitrogen fixation is a well documented phenomenon that must be present in any mangrove ecosystem [42–47]. Several species of N_2 -fixing mangrove root-associated bacteria were isolated and identified [33,37].

Many species of microorganisms can solubilize phosphate under marine environments. Phosphate-solubilizing bacteria can be found in the water column [48], in the rhizosphere of the seaweed *Zostera marina* [49], and associated with white and black mangroves [20]. Their number can be as high as 10^7 cfu g^{-1} sediment [50]. At present, the interactions, if any, between mangrove root-associated N_2 -fixing bacteria and phosphate-solubilizing bacteria and plants are unknown.

We showed that when two species of mangrove-associated bacteria are mixed in vitro in a culture medium or in seawater, they affect each other's metabolism. When grown in culture medium, the two populations, after keeping a steady state for 24 and 72 h, declined probably because of nutrient exhaustion. When the mixture was grown in seawater, this did not happen and the mixture then had an advantage over single cultures. Although one may expect nutrient exhaustion to be more apparent in seawater, seawater would appear to be a better culture medium for this marine diazotrophic bacteria than the N-free medium used in his study.

The two bacterial species, originally isolated from the same tree, mutually affect one another's metabolism in mixed cultures in vitro. Nitrogen fixation increased in the N_2 -fixing bacterium *Phyllobacterium* sp., and phosphate solubilization increased in *B. licheniformis*. The effect of mixed inoculation on black mangrove seedlings was moderate. The average number of true leaves in the seedlings increased as did the incorporation of ^{15}N into the leaves and stem tissues. This indicates a direct transfer of nitrogen from the N_2 -fixing bacterium to the plant, similar to that which occurred when the diazotrophic cyanobacterium *M. chthonoplastes* was inoculated on black mangrove seedlings [10,12]. However, the total nitrogen content of the plant decreased. We might speculate that the supply of nitrogen by the N_2 -fixing bacterium was insufficient to keep pace with the increased growth of the seedling (from 9.5 to 10 cm tall in 26 days) and to

supply the requirements of the phosphate-solubilizing *B. licheniformis*.

In sum, we showed that bacterial species, each with a potential as a PGPB, can interact when mixed and their metabolism is affected. Though the black mangrove rhizosphere and aerial roots are heavily colonized by numerous bacteria and other protista [37], it is premature to speculate whether the effects that were detected also occur in the rhizosphere, and if so at what magnitude.

Acknowledgements

Y.B. participated in this study in memory of the late Mr. Avner and Mr. Uzi Bashan from Israel. We thank Mr. Alejandro Amador for his technical help in isolating the bacterial strains, Miss Patricia Vazquez, Mr. Juan-Pablo Hernandez, Mr. Manuel Moreno, and Mrs. Esther Puente for valuable technical comments and laboratory help, Mr. Ariel Cruz for gas chromatography, Mrs. Sonia Rocha for nitrogen analysis, Mr. Bill Mark for ^{15}N analysis, and Dr. Ellis Glazier for editing the English text. This study was supported by Consejo Nacional de Ciencia y Tecnología (CONACyT), Mexico, contracts # 26262-B and # 28362-B, by the Bashan Foundation (to Y.B.) and by the Natural Science and Engineering Research Council of Canada (to B.R.G.).

References

- Alongi, D.M. (1994) The role of bacteria in nutrient recycling in tropical mangrove and other coastal benthic ecosystems. *Hydrobiologia* 285, 19–32.
- Alongi, D.M., Boto, K.G. and Tirendi, F. (1989) Effect of exported mangrove litter on bacterial productivity and dissolved organic carbon fluxes in adjacent tropical nearshore sediments. *Mar. Ecol. Prog. Ser.* 56, 133–144.
- Bano, N., Nisa, M.U., Khan, N., Saleem, M., Harrison, P.J., Ahmed, S.I. and Azam, F. (1997) Significance of bacteria in the flux of organic matter in the tidal creeks of the mangrove ecosystem of the Indus river delta, Pakistan. *Mar. Ecol. Prog. Ser.* 157, 1–12.
- Holguin, G. and Bashan, Y. (2001) La importancia de los manglares y su microbiología para el sostenimiento de las pesquerías costeras. In: *Agricultural Microbiology for the 21st Century* (Ferrera-Cerrato, R. and Alarcon, A., Eds.), Published by Mundi Prensas, Mexico City (in press).
- Holguin, G., Bashan, Y., Mendoza-Salgado, R.A., Amador, E., Toledo, G., Vazquez, P. and Amador, A. (1999) La microbiología de los manglares, bosques en la frontera entre el mar y la tierra. *Cienc. Desarrollo* 25 (144), 26–35.
- Holguin, G., Vazquez, P. and Bashan, Y. (2001) The role of sediment microorganisms in the productivity, conservation, and rehabilitation of mangrove ecosystems: An overview. *Biol. Fertil. Soils* (in press).
- Bunt, J.S. (1992) Introduction. In: *Tropical Mangrove Ecosystems* (Robertson, A.I. and Alongi, D.M., Eds.), pp. 1–6. American Geophysical Union, Washington, DC.
- Lacerda, L.D., Conde, J.E., Bacon, P.R., Alarcon, C., Alvarez-Leon, R., D'Croz, L.D., Kjerfve, B., Polania, J. and Vannucci, M. (1993) Ecosistemas de manglar de America Latina y el Caribe: sinopsis. In: *Conservación y Aprovechamiento Sostenible de Bosques de Manglar en las Regiones America Latina y Africa* (Lacerda, L.D. and Polania, J., Eds.), pp. 1–38. International Tropical Timber Organization and International Society for mangrove Ecosystems, Okinawa.
- Twilley, R.R., Pozo, M., Garcia, V.H., Rivera-Monroy, V.H., Zambrano, R. and Boderó, A. (1997) Litter dynamics in riverine mangrove forests in the Guayas River Estuary, Ecuador. *Oecologia* 111, 109–122.
- Bashan, Y., Puente, M.E., Myrold, D.D. and Toledo, G. (1998) In vitro transfer of fixed nitrogen from diazotrophic filamentous cyanobacteria to black mangrove seedlings. *FEMS Microbiol. Ecol.* 26, 165–170.
- Puente, M.E., Holguin, G., Glick, B.R. and Bashan, Y. (1999) Root-surface colonization of black mangrove seedlings by *Azospirillum halopraeferens* and *Azospirillum brasilense* in seawater. *FEMS Microbiol. Ecol.* 29, 283–292.
- Toledo, G., Bashan, Y. and Soeldner, A. (1995) In vitro colonization and increase in nitrogen fixation of seedling roots of black mangrove inoculated by a filamentous cyanobacteria. *Can. J. Microbiol.* 41, 1012–1020.
- Bashan, Y. and Holguin, G. (1997) *Azospirillum*–plant relationships: environmental and physiological advances (1990–1996). *Can. J. Microbiol.* 43, 103–121.
- Glick, B.R. (1995) The enhancement of plant growth by free-living bacteria. *Can. J. Microbiol.* 41, 109–117.
- Glick, B.R., Patten, C.L., Holguin, G. and Penrose, D.M. (1999) *Biochemical and Genetic Mechanisms used by Plant Growth-Promoting Bacteria*. Imperial College Press, London.
- Hallmann, J., Quadt-Hallmann, A., Mahafee, W.F. and Kloepper, J.W. (1997) Bacterial endophytes in agricultural crops. *Can. J. Microbiol.* 43, 895–914.
- Chanway, C.P. and Holl, H.B. (1992) Influence of soil biota on Douglas-fir (*Pseudotsuga menziesii*) seedling growth: the role of rhizosphere bacteria. *Can. J. Bot.* 70, 1025–1031.
- Reinhold, B., Hurek, T., Fendrik, I., Pot, B., Gillis, M., Kersters, K., Thielemans, S. and De Ley, J. (1987) *Azospirillum halopraeferens* sp. nov., a nitrogen-fixing organism associated with roots of Kallar grass (*Leptochloa fusca*) (L.) Kunth. *Int. J. Syst. Bacteriol.* 37, 43–51.
- Holguin, G. and Bashan, Y. (1996) Nitrogen-fixation by *Azospirillum brasilense* Cd is promoted when co-cultured with a mangrove rhizosphere bacterium (*Staphylococcus* sp.). *Soil Biol. Biochem.* 28, 1651–1660.
- Vazquez, P., Holguin, G., Puente, M.E., Lopez-Cortes, A. and Bashan, Y. (2000) Phosphate-solubilizing microorganisms associated with the rhizosphere of mangroves growing in a semiarid coastal lagoon. *Biol. Fertil. Soils* 30, 460–468.
- Bashan, Y., Moreno, M. and Troyo, E. (2000) Growth promotion of the seawater-irrigated oilseed halophyte *Salicornia bigelovii* inoculated with mangrove rhizosphere bacteria and halotolerant *Azospirillum* spp. *Biol. Fertil. Soils* 32, 265–272.
- Alagawadi, A.R. and Gaur, A.C. (1992) Inoculation of *Azospirillum brasilense* and phosphate-solubilizing bacteria on yield of (*Sorghum bicolor* (L.) Moench) in dry land. *Trop. Agric.* 69, 347–350.
- Belimov, A.A., Kojemiakov, P.A. and Chubarliyeva, C.V. (1995) Interaction between barley and mixed cultures of nitrogen fixing and phosphate-solubilizing bacteria. *Plant Soil* 173, 29–37.
- Del Gallo, M. and Fabbri, P. (1991) Effect of soil organic matter on chickpea inoculated with *Azospirillum brasilense* and *Rhizobium leguminosarum* bv. *ciceri*. *Plant Soil* 137, 171–175.
- Elshanshoury, A.R. (1995) Interactions of *Azotobacter chroococcum*, *Azospirillum brasilense* and *Streptomyces mutabilis*, in relation to their effect on wheat development. *J. Agron. Crop Sci.* 175, 119–127.
- Flouri, F., Sini, K. and Balis, C. (1995) Interactions between *Azospirillum* and *Phialophora radicola*. *NATO ASI Ser. G-37*, 231–237.
- Hassouna, M.G., Hassan, M.T. and Madkour, M.A. (1994) Increased yields of alfalfa (*Medicago sativa*) inoculated with N_2 -fixing bacteria and cultivated in a calcareous soil of Northwestern Egypt. *Arid Soil Res. Rehabil.* 8, 389–393.

- [28] Navale, A.M., Shinde, D.B., Vaidya, B.R. and Jadhav, S.B. (1995) Effect of *Azotobacter* and *Azospirillum* inoculation under graded levels of nitrogen on growth and yield of sugarcane (*Saccharum officinarum*). Indian J. Agron. 40, 665–669.
- [29] Drozdowicz, A. and Ferreira Santos, G.M. (1987) Nitrogenase activity in mixed cultures of *Azospirillum* with other bacteria. Zent.bl. Mikrobiol. 142, 487–493.
- [30] Isopi, R., Fabbri, P., Del Gallo, M. and Puppi, G. (1995) Dual inoculation of *Sorghum bicolor* (L.) Moench ssp. *bicolor* with vesicular arbuscular mycorrhizas and *Acetobacter diazotrophicus*. Symbiosis 18, 43–55.
- [31] Halsall, D.M. and Gibson, A.H. (1989) Nitrogenase activity of a range of diazotrophic bacteria on straw, straw breakdown products and related compounds. Soil Biol. Biochem. 21 (2), 291–298.
- [32] Khammas, K.M. and Kaiser, P. (1992) Pectin decomposition and associated nitrogen fixation by mixed cultures of *Azospirillum* and *Bacillus* species. Can. J. Microbiol. 38, 794–797.
- [33] Holguin, G., Guzman, M.A. and Bashan, Y. (1992) Two new nitrogen-fixing bacteria from the rhizosphere of mangrove trees, isolation, identification and in vitro interaction with rhizosphere *Staphylococcus* sp.. FEMS Microbiol. Ecol. 101, 207–216.
- [34] Knösel, D.H. (1984) Genus *Phyllobacterium*. In: Bergie's Manual of Systematic Bacteriology, Vol. 1 (Kreig, N.R. and Holt, J.G., Eds.), pp. 254–256. The Williams and Wilkins Co., Baltimore, MD.
- [35] Lambert, B., Joos, H., Dierickx, S., Vantomme, R., Swings, J., Kersters, K. and Van Montagu, M. (1990) Identification and plant interaction of a *Phyllobacterium* sp., a predominant rhizobacterium of young sugar beet plants. Appl. Environ. Microbiol. 56, 1093–1102.
- [36] Giani, L., Bashan, Y., Holguin, G. and Strangmann, A. (1996) Characteristics and methanogenesis of the Balandra lagoon mangrove soils, Baja California Sur, Mexico. Geoderma 72, 149–160.
- [37] Toledo, G., Bashan, Y. and Soeldner, A. (1995) Cyanobacteria and black mangroves in northwestern Mexico: colonization, and diurnal and seasonal nitrogen fixation on aerial roots. Can. J. Microbiol. 41, 999–1011.
- [38] Postgate, J.R. (1969) Viable counts and viability. Methods Microbiol. 1, 611–627.
- [39] Carley, H.E. and Watson, R.D. (1966) A new gravimetric method for estimating root-surface areas. Soil Sci. 102, 289–291.
- [40] Barrie, A. and Prosser, S.J. (1996) Automated analysis of light-element stable isotopes ratio mass spectrometry. In: Mass Spectrometry of Soils (Boutton, T.W. and Yamasaki, S., Eds.), pp. 1–46. Marcel Dekker, New York.
- [41] APHA-AWWA-WPCF (1989) Standard methods for the examination of water and wastewater. 17th edn., American Public Health Association, Washington, DC.
- [42] Hicks, B.J. and Silvester, W.B. (1985) Nitrogen fixation associated with the New Zealand mangrove (*Avicennia marina* (Forsk.) Vierh. var. *resinifera* (Forst. F) Bakh). Appl. Environ. Microbiol. 49, 955–959.
- [43] Jones, K. (1992) Diurnal nitrogen fixation in tropical marine cyanobacteria: a comparison between adjacent communities of non-heterocystous *Lyngbya* sp. and heterocystous *Calothrix* sp.. Br. Phycol. J. 27, 107–118.
- [44] Potts, M. (1979) Nitrogen fixation (acetylene reduction) associated with communities of heterocystous and non-heterocystous blue-green algae in mangrove forests of Sinai. Oecologia 39, 359–373.
- [45] van der Valk, A.G. and Attiwill, P.M. (1984) Acetylene reduction in an *Avicennia marina* community in Southern Australia. Aust. J. Bot. 32, 157–164.
- [46] Zuberer, D.A. and Silver, W.S. (1978) Biological dinitrogen fixation (acetylene reduction) associated with Florida mangroves. Appl. Environ. Microbiol. 35, 567–575.
- [47] Zuberer, D.A. and Silver, W.S. (1979) N₂-fixation (acetylene reduction) and the microbial colonization of mangrove roots. New Phytol. 82, 467–471.
- [48] Promod, K.C. and Dhevendaran, K. (1987) Studies on phosphobacteria in Cochin backwater. J. Mar. Biol. Assoc. India 29, 297–305.
- [49] Craven, P.A. and Hayasaka, S.S. (1982) Inorganic phosphate solubilization by rhizosphere bacteria in a *Zostera marina* community. Can. J. Microbiol. 28, 605–610.
- [50] Devenderan, K., Sundararaj, V., Chandramohan, D. and Krishnamurthy, K. (1974) Bacteria and primary production. Indian J. Mar. Sci. 3, 139–141.